# VSV-G Tag Antibody

Goat Polyclonal

Antigen Affinity Purified

Catalog No. A190-130A

Lot No. 260817

| | |
|---|---|
| **APPLICATIONS** | WB, IP, ICC, ELISA |
| **AMOUNT** | 0.1 ml |
| **CONCENTRATION** | 1 mg/ml |
| **STORAGE/SHELF LIFE** | 2 - 8°C / 1 year from date of receipt |
| **PHYSICAL STATE** | Liquid |
| **BUFFER** | Phosphate Buffered Saline (PBS) containing 0.09% Sodium Azide |
| **ISOTYPE** | IgG |
| **ORIGIN** | USA |
| **PRODUCTION PROCEDURES** | Goats were immunized with VSV-G (YTDIEMNRLGK) conjugated to KLH. Antibody was isolated by affinity chromatography using the peptide immobilized on solid support.<br><br>Prior to conjugation, immunoglobulin concentration was determined using Beer's Law where 1mg/mL IgG has an A280 of 1.4. |
| **APPLICATIONS** | Centrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use. |

| | |
|---|---|
| Western Blot | 1:1000 - 1:30,000 |
| Immunoprecipitation | 1 - 4 µg/mg lysate |
| Immunocytochemistry | 1:100 - 1:400 |
| ELISA | 1:1000 - 1:30,000; for coating plates 1:100 - 1:500 |


| | |
|---|---|
| **APPLICATION NOTES** | All western blot analysis is performed using 5% Milk-TBST for blocking and as antibody diluent. Primary antibody is incubated overnight.<br><br>Western blots are performed using Donkey anti-Goat IgG Heavy and Light Chain Antibody (A50-101P). |
| **ADDITIONAL INFO** | Please visit our website for additional product information. |

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.

Date: August 17, 2026



For in vitro laboratory use only. Not for any clinical, therapeutic or diagnostic use in humans or animals. Not for human or animal consumption. This product may not be resold or modified for resale without the prior written approval of Bethyl Laboratories, Inc. The information provided in this data sheet is believed to be correct but does not purport to be all-inclusive and is intended to be used as a guide. Bethyl Laboratories, Inc. shall not be liable or responsible in any way for use of either this information or the material supplied. Disposal of hazardous material may be subject to federal, state or local laws or regulations.




**Detection of VSV-G-tagged protein by western blot.** *Samples:* 200, 100, or 50 ng of E. coli whole cell lysate expressing a multi-tag fusion protein. *Antibodies:* Affinity purified, goat anti-VSV-G antibody A190-130A used for WB at 0.04 µg/ml (1:25,000). *Detection:* Chemiluminescence with an exposure time of 30 seconds.



For in vitro laboratory use only. Not for any clinical, therapeutic or diagnostic use in humans or animals. Not for human or animal consumption. This product may not be resold or modified for resale without the prior written approval of Bethyl Laboratories, Inc. The information provided in this data sheet is believed to be correct but does not purport to be all-inclusive and is intended to be used as a guide. Bethyl Laboratories, Inc. shall not be liable or responsible in any way for use of either this information or the material supplied. Disposal of hazardous material may be subject to federal, state or local laws or regulations.