

CTLA4 Recombinant Monoclonal Antibody [BLR257L]

Rabbit Recombinant Monoclonal

Purified		RefSeq ID	NP_005205.2
Catalog No.	A700-257	Uniprot ID	P16410
Lot No.	1	GeneID	1493

APPLICATIONS	WB, IHC, mIF						
SPECIES REACTIVITY	Human						
AMOUNT	100 µl						
CONCENTRATION	100 µg/ml						
STORAGE/SHELF LIFE	2 – 8°C / 1 year from date of receipt						
PHYSICAL STATE	Liquid						
BUFFER	Borate Buffered Saline (BBS) pH 8.2 with 0.1% BSA and 0.09% Sodium Azide						
ISOTYPE	IgG						
CLONE #	BLR257L						
ORIGIN	USA						
PRODUCTION PROCEDURES	<p>Recombinant antibody was purified from cell culture supernatant.</p> <p>Immunogen was a peptide representing a region between residue 173 and the C-terminus (residue 223) of human Cytotoxic T-lymphocyte protein 4 using the numbering given in entry NP_005205.2 (Gene ID 1493).</p>						
APPLICATIONS	<p>Centrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use.</p> <table><tr><td>Western Blot</td><td>1:1,000</td></tr><tr><td>Immunohistochemistry</td><td>1:100 to 1:500. Epitope retrieval with citrate buffer pH 6.0 is recommended for FFPE tissue sections.</td></tr><tr><td>Multiplex Immunofluorescence</td><td>1:20 (COMET™)</td></tr></table>	Western Blot	1:1,000	Immunohistochemistry	1:100 to 1:500. Epitope retrieval with citrate buffer pH 6.0 is recommended for FFPE tissue sections.	Multiplex Immunofluorescence	1:20 (COMET™)
Western Blot	1:1,000						
Immunohistochemistry	1:100 to 1:500. Epitope retrieval with citrate buffer pH 6.0 is recommended for FFPE tissue sections.						
Multiplex Immunofluorescence	1:20 (COMET™)						
IHC HUMAN CONTROLS	Appendix, Lymph Node, Spleen, Tonsil						
ADDITIONAL INFO	Please visit our website for additional product information.						

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.
Michael Spencer, PhD Date: July 3, 2023